

# GET IN THE ZONE WITH “ZONE 1”

## For Use by Licensed Professionals Only.

This “All Purpose” 5% lidocaine topical anesthetic has the advantage of a pH of 7, for rapid onset and safe use around the eyes- although care should be taken to avoid getting this product in the eyes. The cream base also makes this product self-occlusive. This product is for relief of discomfort due to anorectal disorders, or in preparation for other pain sensitive procedures such as large tattooing.

**FOR EXTERNAL USE ONLY ON INTACT SKIN:** Check with client for history of allergic reactions to topical anesthetics before applying. Keep in a cool dark place, *out of the reach of children.*

**WARNING:** Avoid direct contact with the eyeball. If contact should occur, irrigate with copious amounts of sterile saline and contact an eye care physician. **DO NOT SWALLOW.** If accidentally swallowed, seek medical help immediately.

SUGGESTED APPLICATION TIMES

### EYEBROWS

15-20 minutes

We suggest using *Pro Plus* for this procedure. Use *SJ3* during the procedure to continue anesthesia, and either *Zone 2* or *Blue Gel* to continue anesthesia and constrict blood vessels.

### EYELINER

10- 15minutes

Fill the eyes with liquid eye gel to coat the cornea. With the client's eyes closed, apply a thin layer of cream with a small applicator across the lash line. Remove anesthetic before beginning procedure. Following the first pass of tattooing, sparingly apply *Zone 2* or *Blue Gel* to control swelling and maintain anesthesia for up to two hours.

### LIPS

15-20 minutes

We suggest using *Pro Plus* or *Super T* for this procedure. While some relief may be provided with topical anesthetics, for total anesthesia, you must provide a nerve block performed by a physician. *Do not use inside of the mouth.* Use *SJ3* during procedure to continue anesthesia, or either *Blue Gel*, *Vasocaine* or *Zone 2* to continue anesthesia and constrict blood vessels.

### BODY SKIN

90-120 minutes

Prior to tattooing or piercing: Lightly exfoliate with *Skin Resurfacer*. Gently massage into skin and then apply a medium coating over the top. In order to prevent absorption into clothing and retain effective anesthesia, cover with an occlusive dressing. Thicker skin requires longer time to absorb the cream than thinner skin. Use *SJ3* during procedure to continue anesthesia, or either *Blue Gel*, *Vasocaine* or *Zone 2* to continue anesthesia and constrict blood vessels.

### PIERCING

15- 20 minutes

We suggest using *Super T* for this procedure. Avoid genital areas. Thicker skin requires longer pre-treating time. Lightly exfoliate with *Skin Resurfacer*. Gently massage into skin, then apply a medium coating on top. Wipe away immediately prior to piercing. Use *Blue Gel* or *Zone 2* during jewelry insertion to continue anesthesia and reduce swelling.

**DERMAL SOURCE**  
1017 SW Morrison Street, #210  
Portland, Oregon 97205  
Phone: 866-568-3223, Fax: 503-973-5433  
[www.dermalsouce.com](http://www.dermalsouce.com)

